

EFFECT OF PROTEIN LEVEL AND HAPA EXCHANGE TIMES ON GROWTH PERFORMANCE AND PROFITABILITY OF NILE TILAPIA (*OREOCHROMIS NILOTICUS*) FRY DURING SEX REVERSAL STAGE IN COMMERCIAL HATCHERY, FAYOUM GOVERNORATE, EGYPT

R.M. Abou Zied

Faculty of Agriculture, Fayoum University, Fayoum, Egypt.

(Received 9/6/2015, accepted 22/7/2015)

SUMMARY

Two trials were conducted for three weeks (one during June and the other during July 2014) to evaluate the effect of protein level and hapa exchange times on growth performance and profitability of Nile tilapia (*Oreochromis niloticus*) fries during sex reversal stage in a commercial hatchery, Fayoum governorate, Egypt. The first one was conducted in six hapa (2 per each treatment) with size of 10.2 m³ (4 × 3 × 0.85 m). The hapas were stocked with Nile tilapia fry at a rate of 50,000 fry/hapa. Three diets varying in protein level [fish meal as animal protein source] were formed as follows: the D₇₂ [100% fish meal (FM) = 72 % Crude Protein (CP)]; the D₅₃ (70% FM + 30% wheat flour = 53% CP) and D₃₇ (33.33% FM + 33.33 % wheat flour + 33.33 % commercial diets = 37% CP). The diets were offered to the fry five times/day at a rate of 30% of total biomass weight/first week then decreased gradually by 10% every week. The second trial was to study the effect of hapa exchange times [zero time (OTT); one time (1TT) and two times (2TT) treatments, where, every treatment had two replicates] on the fry growth performance. The fry were fed with D₃₇ (37% CP) which was mentioned previously at the first trail at the same feeding rate and times/day. The hapas of both first and second trails were fixed in an earthen pond (3000 m²).

The results obtained from the first trail revealed that significant differences ($P \leq 0.05$) were found for all growth performance parameters. The D₇₂ has the best values for final weight, weight gain, daily gain, specific growth rate. On the other hand, survival rate and feed conversion ratio have an opposite trend for that treatment (D₇₂) when it was compared with the other treatments. But the profit index (PI) was significantly higher ($P \leq 0.05$) for fish fed D₃₇ followed by D₅₃ and finally D₇₂.

The results of the second trail demonstrated that there were significant differences ($P \leq 0.05$) for all growth parameters the best values was obtained with both 1TT and 2TT when were compared with OTT.

Keywords: *Nile tilapia (O. niloticus), fish meal, protein level, hapa exchanged growth performance, profit index.*

INTRODUCTION

Tilapia is the second aquaculture species globally. Egypt occupies the second ranked in the production of tilapia after China (FAO, 2014), because of it's easy to adapt in tropical and sub-tropical areas of the world (Shelton, 2002). Tilapia have numerous advantages as an aquaculture species (Teichert-Coddington *et al.*, 1997) but the ability to reproduce in the production setting has resulted in various techniques being developed to control unwanted reproduction.

Net enclosures or hapas are used in tilapia hatcheries (Guerrero, 1997). Hapas has many attributes that make them an excellent hatchery system for tilapia, especially in developing countries (El-Sayed, 2006). These include easy construction, easy management, easy seed harvest and low cost. Hapas can also be suspended in fertilized earthen ponds, deep water bodies and concrete tanks supplied with clear water.

As the main and most expensive component of the diet, protein draws greater attention in nutrition requirement studies (Loum *et al.*, 2013).

Fry feed generally contains higher level of protein because protein and energy requirements are higher in the early stages of life. Protein content of fry feed for tilapia farming has not been standardized yet although some farms use feed stuffs having 40% protein. So, it is essential to recommend the appropriate protein level of fry feed for economic production of healthy fry and maximize its lifespan as well (Loum *et al.*, 2013). So, the amount of protein in the diet should be just enough for fish growth where the excess protein in fish diets may be wasteful and cause diets to be unnecessarily expensive (Ahmad, 2000). Reducing feeding costs could be a key factor for successful development of aquaculture.

Fry growth is improved with increasing protein levels. Larvae of tilapia tend toward carnivores, yet as they mature they become more herbivorous (Suresh, 2003). Thus, dietary formulations may require modification as fish grow and ingredient usage may shift from animal-based ingredients such as fish meal to plant-based feedstuffs. Most early life-history stage diets for tilapia contain higher concentrations of fish meal than grow out diets (Gonzales *et al.*, 2007).

Klanian and Adame (2013) reported that water was changed with new water, to dilute the high levels of ammonia and nitrites. Water exchange and flow rate may affect the growth and physiological functions of tilapia. Little information's are available on the subject. Continuous water exchange generally sustains the good quality of culture water while low or zero water exchange may reduce the quality of water lead to reduce growth and increased mortality (El-Sayed *et al.* 2005).

This work aimed to study the effects of protein level and hapa exchange times on growth performance, survival rate and profitability of Nile tilapia (*O. niloticus*) fry during sex reversal stage in commercial hatcheries.

MATERIALS AND METHODS

Two experiments were conducted at commercial hatchery in Shakshouk village, Fayoum governorate, Egypt for three weeks during summer season 2014 (the first during June and the second during July) to evaluate the effect of protein level and hapa exchange times on growth performance and profitability of Nile tilapia (*O. niloticus*) fry. After hatching and before consumption of yolk sac, the fry were collected, graded, selected and weighed to be prepare for the study.

The first experiment was conducted in six hapa (two hapas per each treatment) with size of 10.2 m³ (4 × 3 × 0.85 m). They were stocked with Nile tilapia fry at a rate of 50,000 fry/hapa. The fry were fed with three diets [72%, 53% and 37% crude protein (CP)] as follows: the D₇₂ [100% fish meal (FM) = 72% CP]; the D₅₃ (70% FM + 30% wheat flour = 53% CP) and the D₃₇ (33.33% FM + 33.33 % wheat flour + 33.33 % commercial diets =37% CP). The diets were treating with 17 α methyl testosterone hormone at a rate of 100 mg /kg diet (Al-Hakim *et al.*, 2012). The diets were applied to the fry at a rate of 30% of total biomass, where the daily quantities were divided to five portions/day. After the first week, the quantities were decreased gradually by 10% every week (three weeks experimental period).

In the second experiment the fry were stocked inside the hapas (with 10.2 m³) in an earthen pond (0.75 feddan = 3000 m²) with different exchange times (two replicates per each). Firstly, zero time treatment (0TT), where, 50,000 fry were stocked at hapa (10.2 m³) from the first day till 21 days without any exchange. Secondly, one time treatment (1TT), where 50,000 fry were stocked at hapa (10.2 m³) from the first day until 10 days, then the fry were transferred in a new hapa (10.2 m³) till 21 days. And finally, two times treatment (2TT), where 50,000 fry were stocked at hapa (10.2 m³) after 7 and 15 days to the end of the trail (21 days). Fry were fed with D₃₇ (37% CP) plus hormone incorporated feed five times/day. Weekly samplings of both water and fry were done to test the water quality parameters and to determine the growth performance characteristics and profitability. All treatments were started during the first 48th h, this due to the hatchery production of fry was not enough to start all treatment in the same time. A plan of the second trail is shown in Table (1).

Parameters measurements:

Some water quality parameters were recorded weekly during the study period for the two trails. Water temperature, pH, dissolved oxygen (DO), ammonia nitrogen (NH₃-N), and salinity were measured weekly by

centigrade thermometer, Orion digital pH meter model 201, oxygen meter, Cole Parmer model 5946, HACH test kit ammonia mid-range 0-3 mg/L model NI-8, and TDS apparatus, respectively.

Table (1). A plan of the second trail.

Item	Sex reversal stage in days																				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
0TT	50,000 fry/ 10.2 m ³																				
1TT	50,000 fry/10.2 m ³										→ 50,000 fry/10.2 m ³										
2TT	50,000 fry/10.2 m ³							→ 50,000 fry/10.2 m ³							→ 50,000 fry/10.2 m ³						

Note that:

- 0TT = zero time treatment (the hapa was not exchanged);
- 1TT= one time treatment (the hapa was exchanged only one time) and
- 2TT= two time treatment (the hapa was exchanged two times).

By the end of both trails, some growth parameters and survival rate were measured

- Weight gain = Final weight - Initial weight (Effiong *et al.*, 2009),
- Daily gain = Weight gain, g /period in days (Effiong *et al.*, 2009),
- Specific growth rate (SGR %) = 100 (ln Final weight-ln Initial weight)/period in days, where ln is the natural log (Effiong *et al.*, 2009),
- Feed conversion ratio (FCR) = feed offered/weight gain (Effiong *et al.*, 2009),
- Survival rate (SR) % = Final number of fish /Initial number of fish x 100 (Charo-Karisa *et al.*, 2006) and
- Condition factor (CF) =100 x W/L³ where W= Weight (g), L = Total length (cm) (Ng and Wang, 2011).

Economic analysis:

A simple economic analysis was used to assess the cost effectiveness of diets used in the feed trial. The cost of feed was calculated using market prices, taking into consideration the cost of feed and the transport fare with the assumption that all other operating costs remained constant (e.g. cost of constructing hapa, cost of broodstock and labor).

Indices for economic evaluation included:

(i) Incidence cost (IC), which was calculated as:

$$IC = \text{Cost of feed} / \text{No. of fry produced}$$

IC is actually the cost of feed to produced 1000 fry (relative cost per unit), and the lower the value, the more profitable using that particular feed (Nwanna, 2003; Abu *et al.*, 2010).

(ii) Profit index (PI), which was calculated as:

$$PI = \text{value of fish produced} / \text{Cost of feed}$$

Statistical Analysis:

Data were computed and statistically analyzed using a one-way analysis of variance using SPSS version 16 (2007). Mean of treatments were compared by Duncan multiple range test when the differences were significant (Duncan 1955).

RESULTS AND DISCUSSION

Water quality parameters:

Some water quality parameters were recorded during the study. The dissolved oxygen (DO) was 5.7 ± 0.14 mg/l. The pH was 8.05 ± 0.13 . The temperature was 29 ± 1.0 °C. The $\text{NH}_3\text{-N}$ concentration was 0.4 ± 0.17 mg/l and salinity 2.8 ± 0.22 ‰. These values were within the safe and acceptable ranges for the spawning and growth of Nile tilapia fry and agreed with that reported by many authors (Magid and Babiker, 1975; Ross, 2000; El-Sayed, 2006 and El-Sherif & El-Feky, 2008).

First experiment: Diets' protein levels:

As shown in Table (2), the D_{72} has highest significant ($P \leq 0.05$) values for all growth performance parameters except survival rate and FCR, where it has the lowest one. This may be due to firstly to: higher protein content of D_{72} and fish meal diets are also assumed to contain androgenic hormones which are beneficial for fish growth, and, secondly to: the androgenic steroids which promote releasing of growth hormone from pituitary somatotrophs fish (Higgs *et al.*, 1976).

Table (2). Effects of diets' protein levels on some parameters of growth performance of Nile tilapia (*O. niloticus*) fry reared in hapa (50,000 fry/10.2 m³) for 21 days.

Item	Initial weight, mg	Final weight, mg	Weight gain, mg	Daily gain, mg	SGR, %	Survival rate, %	FCR
D_{72}	10.99	149.99 ^a	138.00 ^a	6.57 ^a	12.41 ^a	75.55 ^b	1.38 ^b
D_{53}	10.99	101.47 ^b	90.48 ^b	4.31 ^b	10.58 ^b	92.85 ^a	1.72 ^a
D_{37}	10.99	99.87 ^b	88.88 ^b	4.23 ^b	10.51 ^b	94.17 ^a	1.73 ^a
SED	0.01	3.24	3.24	0.15	0.12	2.16	0.04

Note that:

- Values in the same column with different superscripts are significantly different ($P \leq 0.05$).

- SED, standard error of a difference between 2 means = $\sqrt{2 \times \text{Error MS}/r}$

- D_{72} [100% fish meal (FM) = 72% CP]; D_{53} (70% FM + 30% wheat flour = 53% CP) and the D_{37} (33.33% FM + 33.33% wheat flour + 33.33% commercial diets = 37% CP).

D_{37} and D_{53} have highest significant ($P \leq 0.05$) values for both survival rate and FCR. These may be due to a higher content of carbohydrates which lightweight and caused feed mash to float on the water surface. This provide a chance for fry to consume feed, on the other hand for that of D_{72} , where the feed mash were sinking quickly, and so that, all fry were not capable of consuming feed mash at the same time and lead to more mortality.

The trend of the growth of Nile tilapia fry during trial period revealed that tilapia fed with fish meal consistently increases rapidly in terms of body weight followed by those treated with fish meal with flour (70:30) and the fish meal with flour and artificial feed (equally).

Several dietary protein requirements of several tilapia species have been estimated to range between 20-56% (El-Sayed and Teshima, 1991). De Silva and Perera (1985) reported that the optimum dietary protein level for optimum growth of Nile tilapia fry was 30% crude protein. As reported by Ahmad *et al.* (2004), the growth performance of Nile tilapia fry was highest at 45% protein diets. Al-Hafedh (1999) found that better growth of tilapia was obtained at high dietary protein levels (40-46%) rather than 25-35%. In the present study, fish meal diets contain the highest protein level compared to the other diets which contain 53 and 37% crude protein.

El-Sayed and Teshima (1991) reported that the dietary protein requirements of several species of tilapia have been estimated to range from 20 to 56%. Therefore, protein content of the test diet used in this study was within the suitable range for tilapia culture.

Table (3). Effects of diets' protein levels on profitability of Nile tilapia (*O. niloticus*) fry reared in hapa (50,000 fry/10.2 m³) for 21 days.

Treatment	Feed offered, Kg	Price of hormone, L.E	Feed cost, L.E	Fry No sales	Values of fry sales, L.E	Profit index	Incidence cost
D ₇₂	7250	26.1	131	38000 ^b	950 ^b	6.07 ^c	3.45
D ₅₃	7250	26.1	98	46650 ^a	1166 ^a	9.41 ^b	2.10
D ₃₇	7250	26.1	62	47150 ^a	1179 ^a	13.44 ^a	1.31
SED	--	--	--	867	21.68	0.18	0.13

Note that:

- Values in the same column with different superscripts are significantly different ($P \leq 0.05$).

- SED, standard error of a difference between 2 means = $\sqrt{2 \times \text{Error MS}/r}$

- D₇₂ [100% fish meal (FM) = 72% CP]; D₅₃ (70% FM + 30% wheat flour = 53% CP) and the D₃₇ (33.33% FM + 33.33% wheat flour + 33.33% commercial diets = 37% CP).

As illustrated in Table (3), the cost per kilogram of feed was highest for diet contained 100% fish meal (D₇₂) and lowest for D₃₇. D₃₇ has the highest Survival rate with estimated value of 1179 L.E, but D₇₂ has the lowest fry fed with estimated value of 950 L.E. Total harvested fry with estimated values for diets was in the following order: diet D₃₇ > D₅₃ > D₇₂. Among treatments, D₇₂ recorded the highest significant ($P \leq 0.05$) value of cost/1000 fry (4.13 ± 0.29 L.E) followed by D₅₃ (2.66 ± 0.35) and, finally, D₃₇ (1.87 ± 0.36). The profit index (PI) showed that the cost analysis revealed a lowest cost and highest profit index for the D₃₇. Low cost value means low cost to produce 1000 sex reversed fry. Meanwhile, a high profit index indicates a high amount of profit for every cost incurred in feeding. These are important factors for fish farmer minding when feed ingredients were chosen to minimize cost and maximize profit (Teves *et al.*, 2014).

Second experiment (hapa exchange rate):

Fry hapas in commercial hatcheries are not washed or cleaned for fear of increased mortality for that exchange hapa or replaced by another was necessary. Table (4) demonstrated that 1TT and 2TT treatments lead to improve the growth performance and survival rate and consequently increase profit index. There were significant ($P \leq 0.05$) differences for all growth performance parameters and the higher values were obtained from 1TT and 2TT treatments than OTT treatment. These may be due to cleaning mesh nets allowed to renew and pass the water through the nets and thereby increase the dissolved oxygen and decrease the ammonia concentration, which lead to increase growth rates. Moreover, the OTT treatments provided a chance to accumulate and fill inside mesh nets of the hapa by the feces and feed mash wastes. This prevent to renew and pass the water through the nets, so, lack of oxygen inside the hapa caused weakens growth rate and increased mortality.

Table (4). Effect of frequency hapa exchanged on growth performance and profit index of Nile tilapia fry reared in hapa (50,000 fry/10.2 m³) for 21 days.

Item.	Initial weight	Final weight	Weight gain	SGR	Survival rate	FCR	Profit index
OTT	10.99	89.53 ^b	78.55 ^b	9.99 ^b	71.22 ^b	2.75 ^a	9.65
1TT	10.99	107.62 ^a	96.63 ^a	10.86 ^a	83.96 ^a	1.83 ^b	10.84
2TT	10.99	103.45 ^a	92.46 ^a	10.68 ^a	87.00 ^a	1.84 ^b	10.70
SED	0.07	5.97	5.91	0.26	2.62	0.063	1.33

Note that:

- Values in the same column with different superscripts are significantly different ($P \leq 0.05$).

- SED, standard error of a difference between 2 means = $\sqrt{2 \times \text{Error MS}/r}$

- OTT = zero time treatment (the hapa was not changed);

- 1TT = one time treatment (the hapa was changed only one time) and

- 2TT = two time treatment (the hapa was changed two times).

CONCLUSION

The results obtained from the first trail revealed that the D72 has highest significant ($P \leq 0.05$) values for all growth performance parameters except survival rate and FCR when compared with the other treatments. But, the profit index (PI) was significantly higher ($P \leq 0.05$) for D₃₇ followed by D₅₃. The D₃₇ has lowest cost and the highest profit index in. The second trail demonstrated significant ($P \leq 0.05$) differences for all growth parameters and the 1TT recorded the best values of the studied growth parameters. Depending on the above discussion, exchanging hapa with intense care, at least for one time or more, is preferable to improve environment water quality parameters this will reflex on healthy fry with high growth performance parameters.

REFERENCES

- Abu, O.M.G.; L.O. Sanni; E.S. Erondu and O.A. Akinrotimi (2010). Economic viability of replacing maize with whole cassava root meal in the diet of Hybrid Cat-fish. *Agric. J.*, 1: 1-5.
- Ahmad, M.H. (2000). Improve productive performance in fish. Ph.D. Dissertation, Animal Prod. Department, Faculty of Agriculture, Zagazig University, Egypt.
- Ahmad, M.H.; M. Abdel-Tawwab and Y.A.E. Khattab (2004). Effect of dietary protein levels on growth performance and protein utilization in Nile tilapia (*Oreochromis niloticus* L.) with different initial body weights. Proceedings 6th International Symposium on Tilapia in Aquaculture. 12-16 September 2004, Manila, Philippines. Pp 249-263.
- Al-Hafedh, Y.S. (1999). Effects of dietary protein on growth and body composition of Nile tilapia, *Oreochromis niloticus* L. *Aquaculture Research*, 30(5): 385-393.
- Al-Hakim N.F.A.; M. Saleh; A.Z. Hegazi; A. Ibrahim and K. Aly (2012). Induction of mono-sex (male tilapia) population by inter-specific hybridization and hormonal sex reversal of Nile tilapia. *Egypt J Aquat Biol and Fish*, 1: 23-33.
- Charo-Karisa H.; H. Komen; S. Reynolds; M.A. Rezk; R.W. Ponzoni and H. Bovenhuis (2006). Genetic and environmental factors affecting growth of Nile tilapia (*Oreochromis niloticus*) juveniles: Modeling spatial correlations between hapas. *Aquaculture*, 255: 586-596.
- De Silva, S.S. and M.K. Perera (1985). Effects of dietary protein levels on growth, food conversion and protein use in young *Tilapia nilotica* at four salinities. *Transaction of the American Fisheries Society*, 114: 584-589.
- Duncan, D.B. (1955). Multiple range and multiple F tests. *Biometrics* 11: 1-42.
- Effiong B.N.; A. Sanni and J.O. Fakunle (2009). Effect of partial replacement of fishmeal with duckweed (*Lemna paucicostata*) meal on the growth performance of *Heterobranchus longifilis* fingerlings. *Report Opin.*, 1(3):76-81.
- El-Sayed, A.M. and S. Teshima (1991). Tilapia nutrition in aquaculture. *Reviews in Aquatic Sciences*, 5: 24-265.
- El-Sayed, M.A.; M. Kawanna and M. Mudar (2005). Effect of water flow rates on growth and survival of Nile tilapia fry. *World Aquaculture*, 36 (1): 5-6.
- El-Sayed, A.M. (2006). *Tilapia culture*. CAB International, Wallingford, UK. 277 pp.
- El-Sherif, M.S. and A.M. EL-Feky (2008). Effect of ammonia on Nile Tilapia (*O. niloticus*) performance and some hematological and histological measures. Eighth International Symposium on Tilapia in Aquaculture. Cairo, Egypt.
- FAO (2014). Food and Agriculture Organization of the United Nation. The State of World Fisheries and Aquaculture Opportunities and challenges. Rome,

- Gonzales, Jr. J.M.; A.H. Hutson; M.E. Rosinski; Y.V. Wu; T.F. Powless and P.B. Brown (2007). Evaluation of fish meal-free diets for first feeding Nile tilapia, *Oreochromis niloticus*. Journal of Applied Aquaculture, 19 (3): 89-99.
- Guerrero III., R. D. (1997). A Guide to Tilapia Farming. Aquatic Biosystems. Bay, Laguna, Philippines. 70 pp.
- Higgs, D.A.; E.M. Donaldson; H. Dye and J. R. McBride (1976). Influence of bovine growth hormone and L thyroxin on growth, muscle composition and histological structure of the gonads, thyroid, pancreas and pituitary of coho salmon (*Oncorhynchus kisutch*). J. Fish. Res. Board. Can., 33: 1585- 1603.
- Klanian, M.G. and C.A. Adame (2013). Performance of Nile tilapia *Oreochromis niloticus* fingerlings in a hyper-intensive recirculating aquaculture system with low water exchange. Lat. Am. J. Aquat. Res., 41(1): 150-162.
- Loum, A.; M. Sagne; J. Fall; D. Ndong; M. Diouf; A. Sarr and O.T. Thiaw (2013). Effects of dietary protein level on growth performance, carcass composition and survival rate of fry monosex Nile tilapia, *Oreochromis niloticus* reared under re-circulating system. Journal of Biology and Life Science. 4 (2): 13-22.
- Magid, A. and M.M. Babiker (1975). Oxygen consumption and respiratory behaviour of three Nile fishes. Hydrobiologia 46: 359–367.
- Ng, W.K. and Y. Wang (2011). Inclusion of crude palm oil in the broodstock diets of female Nile tilapia, *Oreochromis niloticus*, resulted in enhanced reproductive performance compared to broodfish fed diets with added fish oil or linseed oil. Aquaculture, 314: 122–131.
- Nwanna L.C. (2003). Risk management in aquaculture by controlled feeding regimen. Pak. J. Nutr., 2(6): 324-328.
- Ross, L.G. (2000). Environmental physiology and energetics. pp. 89–128. In: M. C. M. Beveridge and B. J. McAndrew (eds.) Tilapias: Biology and Exploitation, Fish and Fisheries Series 25, Kluwer Academic Publishers, Dordrecht, the Netherlands.
- Shelton, W.L. (2002). Tilapia culture in the 21st century. P.1-20. in: Guerrero, R.D. III and M.R. Guerrero-del Castillo (eds.). Proceedings of the International Forum on Tilapia Farming in the 21st Century (Tilapia Forum 2002), 184p. Philippine Fisheries Association Inc. Los, Banos, Laguna, Philippines.
- SPSS (2007). Statistical Package for Social Science (for Windows). Release 16 Copyright (C), SPSS Inc., Chicago, USA.
- Suresh, V. (2003). Tilapias. Pages 321-345 in IS. Lucas and P.e. Southgate, eds. Aquaculture: Farming Aquatic Animals and Plants. Blackwell Publishing Co., Oxford,
- Teichert-Coddington, D.R.; T.P. Popma and L.L. Lovshin (1997). Attributes of tropical pond cultured fish. Pages 183–198 in H.S. Egna and C.E. Boyd, eds. Dynamics of Pond Aquaculture. CRC Press, Boca Raton, FL, USA.
- Teves, J.F.C.; M.N.M. Fernandez and J.A. Ragaza (2014). Effects of replacing fishmeal with squash seed meal (*Cucurbita maxima*) on performance of juvenile Nile tilapia (*Oreochromis niloticus*). AACL Bioflux 7(2):68-75.

تأثير مستوى البروتين ومعدل تغيير الهابات على مظاهر النمو والربحية لزريعة البلطي النيلي أثناء مرحلة التحول الجنسي في المفرخات التجارية بمحافظة الفيوم

رمضان محمد أبوزيد

قسم الانتاج الحيوانى- كلية الزراعة- جامعة الفيوم- مصر.

أجريت تجربتان لمدة ثلاثة أسابيع الأولى في شهر يونيه والثانية في شهر يوليو ٢٠١٤ لدراسة تأثير مستوى بروتين العليقة ومعدل تغيير الهابات على مظاهر النمو والربحية لزريعة البلطي النيلي أثناء مرحلة التحول الجنسي في المفرخات التجارية بمحافظة الفيوم- مصر. أجريت الأولى في ستة هابات (هابتان/معاملة) بحجم ١٠,٢ م^٢ (٠,٨٥×٣×٤ م) حملت زريعة البلطي بمعدل ٥٠٠٠٠/هابة. كونت ثلاثة علائق تختلف في نسبة البروتين الأولى (١٠٠% مسحوق سمك = ٧٢% بروتين)، الثانية (٧٠% مسحوق سمك + ٣٠% دقيق قمح = ٥٣% بروتين) أما الثالثة (٣٣,٣% مسحوق سمك + ٣٣,٣% دقيق قمح + ٣٣,٣% عليقة تجارية = ٣٧% بروتين). قدمت العلائق الى الزريعة بعد المعاملة الهرمونية بمعدل ١٠٠ ملجم ١٧ الفا ميثيل تستسترون/كجم خمس مرات/يوم بمعدل ٣٠% من وزن الزريعة في الاسبوع الاول ثم تناقصت تدريجيا بمعدل ١٠% كل اسبوع.

التجربة الثانية درست تأثير عدد مرات تغيير الهابات (بدون تغيير، تغيير مرة، تغيير مرتين) كل معاملة بمكررين على مظاهر النمو. ثبتت الهابات في حوض ترابى (٣٠٠٠ م^٢) وغذيت الزريعة بالعليقة الثالثة (٣٧% بروتين) المستخدمة في التجربة الاولى بنفس معدل التغذية وعدد المرات في اليوم.

أظهرت النتائج المتحصل عليها من التجربة الأولى وجود اختلافات معنوية بين كل مقاييس مظاهر النمو وكانت العليقة الأولى هي الأحسن في الوزن النهائى والزيادة في الوزن و الزيادة اليومية ومعدل النمو النسبى أما معدل الاعاشة والتحويل الغذائى اخذ اتجاه معاكس لباقي المعاملات. دليل الربحية كان أعلى معنويا مع الاسماك المغذاة على العليقة الثالثة تلتها العليقة الثانية واخيرا الأولى. تحليل التكاليف أظهر انخفاض مدى التكاليف وارتفاع دليل الربحية مع العليقة الثالثة.

نتائج التجربة الثانية أظهرت وجود اختلافات معنوية بين كل مظاهر النمو وأحسن القيم التي تم الحصول عليها كانت مع الهابات التي تغيرت مرة واحدة او مرتين بالمقارنة بالهابات التي لم تتغير.