

Demonstrating adverse effects of a common food additive (sodium sulphite) on biochemical, cytological and histopathological parameters in tissues of albino Wister rats

إظهار التأثيرات المعاكسة للمضاف الغذائي الشائع (كبريتيت الصوديوم) علي المقاييس الكيميائية الحيوية والخلوية والنسجية في أنسجة الفئران البيضاء

عنوان البحث

Abstract

Introduction: Sulfites have been used as preservatives in food and drugs. The use of sulfite-containing compounds on fruits and vegetables was banned because of potential allergic reactions, and those suffering from asthma are at elevated risk of sulfite sensitivity.

Method: The present study investigated the effect of daily administration of sodium sulfite (Na_2SO_3) on female albino rats at doses of 200, 500, and 1000 ppm for 12 weeks. Impacts on growth, blood picture, genotoxic effect, some biochemical parameters and histopathological characteristics were examined.

Results: The results revealed that Na_2SO_3 causes a significant decrease in body weight, red blood cells (RBC) count, hemoglobin (Hb) concentration, hematocrit (HCT) value, white blood cells (WBC), and glucose level, while there was a significant increase in serum activity of aspartate aminotransferase (AST), alanine aminotransferase (ALT) and alkaline phosphatase (ALP) as well as the serum urea and creatinine levels in the treated rats. Cytogenetic analysis showed various types of chromosomal aberration such as chromatid gap, chromosome ring, chromatid break, stickiness of chromosome, chromatid separation, and centric fusion. Histopathological examination of the experimental animals indicated little sinusoidal dilatation in rats treated with 200 ppm of Na_2SO_3 . Hepatic vacuolation, large sinusoidal dilatation, degenerative changes and cellular congestion were shown in liver of rats treated with 500 and 1000 ppm of Na_2SO_3 when compared to the control group.

Conclusion: Administration of Na_2SO_3 to rats exhibited serious effects on both liver and kidney cells.