

(Considered Single - Shared with others outside the specialization – Published in International Journal).

Improvement of Selected Morphological, Physiological, and Biochemical Parameters of Roselle (*Hibiscus sabdariffa* L.) Grown Under Different Salinity Levels Using Potassium Silicate and *Aloe saponaria* Extract
Plants, 11(4), 497; (2022) <https://doi.org/10.3390/plants11040497>

Alaa Idris Badawy Abou-Sreea¹, Mohamed H. H. Roby², Hayam A. A. Mahdy³, Nasr M. Abdou⁴, Amira M. El-Tahan⁵, Mohamed T. El-Saadony⁶, Khaled A. El-Tarabily⁷, and Fathy M. A. El-Saadony⁸

¹ Horticulture Department, Faculty of Agriculture, Fayoum University, Fayoum 63514, Egypt;

² Department of Food Science and Technology, Faculty of Agriculture, Fayoum University, Fayoum, Egypt.

³ Botany Department, National Research Centre, Dokki, Giza 12622, Egypt;

⁴ Soil and Water Department, Faculty of Agriculture, Fayoum University, Fayoum 63514, Egypt;

⁵ Plant Production Department, Arid Lands Cultivation Research Institute, The City of Scientific Research and Technological Applications, SRTA-City, Borg El Arab, Alexandria 21500, Egypt;

⁶ Department of Agricultural Microbiology, Faculty of Agriculture, Zagazig University, Zagazig 44511, Egypt;

⁷ Department of Biology, College of Science, United Arab Emirates University, United Arab Emirates

⁸ Agricultural Botany Department, Faculty of Agriculture, Zagazig University, Zagazig 44511, Egypt

Article status

Considered Single - Shared with others outside the specialization – Published in International Journal

Impact Factor: **4.658**

Abstract

Two successive field trials were carried out at the experimental farm of the Agriculture Department of Fayoum University, Fayoum, Egypt, to investigate the sole or dual interaction effect of applying a foliar spray of *Aloe saponaria* extract (Ae) or potassium silicate (KSi) on reducing the stressful salinity impacts on the development, yield, and features of roselle (*Hibiscus sabdariffa* L.) plants. Both Ae or KSi were used at three rates: 0% (0 cm³ L⁻¹), 0.5% (5 cm³ L⁻¹), and 1% (10 cm³ L⁻¹) and 0, 30, and 60 g L⁻¹, respectively. Three rates of salinity, measured by the electrical conductivity of a saturated soil extract (ECe), were also used: normal soil (ECe < 4 dS/m) (S₁); moderately-saline soil (ECe: 4–8 dS/m) (S₂); and highly-saline soil (ECe: 8–16 dS/m) (S₃). The lowest level of salinity yielded the highest levels of all traits except for pH, chloride, and sodium. Ae at 0.5% increased the values of total soluble sugars, total free amino acids, potassium, anthocyanin, a single-photon avalanche diode (SPAD), stem diameter, fruit number, and fresh weight, whereas 1% of Ae resulted in the highest plant height, chlorophyll fluorescence (Fv/Fm), performance index, relative water content, membrane stability index, proline, total soluble sugars, and acidity. KSi either at 30 or 60 g L⁻¹ greatly increased these abovementioned attributes. Fruit number and fruit fresh weight per plant also increased significantly with the combination of Ae at 1% and KSi at 30 g L⁻¹ under normal soil conditions.