

(Considered Single - Shared with others outside the specialization – Published in International Journal).

Chemical analysis of aqueous extracts of *Origanum majorana* and *Foeniculum vulgare* and their efficacy on *Blastocystis spp.* Cysts.

Phytomedicine 43, 158–163. <https://doi.org/10.1016/j.phymed.2018.04.017>

Eman M.H. Méabed^a, Nagwa M. El- Sayed^b, Alaa I.B. Abou-Sreea^c, Mohamed H.H. Roby^d

^a Department of Parasitology, Faculty of Medicine, Fayoum University, Fayoum City, Egypt

^b Department of Parasitology, Research Institute of Ophthalmology, Giza, Egypt

^c Department of Horticulture, Faculty of Agriculture, Fayoum University, Egypt

^d Department of Food Science and Technology, Faculty of Agriculture, Fayoum University, Egypt

| | | |
|----------------|---|-----------------------------|
| Article status | Considered Single - Shared with another outside the specialization – Published in International | Impact Factor: 6.656 |
|----------------|---|-----------------------------|

Abstract

Background: *Origanum majorana* (*O. majorana*) and *Foeniculum vulgare* (*F. vulgare*) are traditionally used herbs in Egypt for treatment of several diseases including parasitic diseases. The Purpose was to determine the efficacy of *O. majorana* and *F. vulgare* aqueous extracts (AEs) on *Blastocystis spp.* in vitro, and to reveal their phenolic, flavonoids components and antioxidant activities through chemical analysis.

Methods: The Efficacy of both plant AEs on human Peripheral Blood Mononuclear Cells (PBMCs) viability was assessed using MTT assay. Isolated *Blastocystis spp.* cysts from patients' diarrhea samples were incubated with different concentrations of *O. majorana* and *F. vulgare* AEs for different incubation periods (24, 48 and 72 h) in comparison with nitazoxanide (NTZ) as a drug control. The total contents of phenolic and flavonoid compounds in the AEs and their ability to reduce DPPH were assessed. High performance liquid chromatography (HPLC) analysis for quantitative and qualitative determination of the phenolic and flavonoid contents was performed. **Results:** *O. majorana* AE at a dose of 400 µg/ml showed efficacy rates of 96% and 100% against *Blastocystis* parasite after 48 and 72 h, respectively, which nearly equivalent to NTZ at a dose of 500 µg/ml. *F. vulgare* at a dose of 250 µg/ml showed less efficacy rate of 56.4% after 48 h and increased to 70.7% after 72 h. Both extracts contain high phenolic and flavonoid compounds that possess antioxidant and free radical scavenging activities.

Conclusion: *O. majorana* and *F. vulgare* AEs showed dose and time dependent anti-Blastocystis activity.