

## conclusion

---

**In conclusion**, the addition of 1 µg/kg of clonidine as an adjuvant to 2% lidocaine and 0.5% bupivacaine in peribulbar anesthesia for cataract surgery significantly decreases the IOP and significantly increases the duration of peribulbar block and improves the analgesic duration without significant side effects. However, clinical trials with a larger number of patients are required to make final conclusions.

### **References:**

- (1) Balbir Khan, Sukhminder Jit Singh Bajwa, Ravi Vohra, Sukhwinder Singh, Rajwinder Kaur, Vartika, and Asha Comparative evaluation of ropivacaine and lignocaine with ropivacaine, lignocaine and clonidine combination during peribulbar anaesthesia for phacoemulsification cataract surgery Indian J Anaesth. 2012 Jan;56(1):21-6
- (2) Neveen Girgis Fahmy MD; Adel Mikhail Fahmy MD, Kareem Yousef Kamal MD. Effect of adding fentanyl and/or clonidine to local anesthetic on prolongation of the peribulbar block in cataract surgery Ain shams journal of anesthesia 2011 Oct volume 4-3
- (3) Hala M. Bahy Eldeen, Mona R. Faheem, Dalia Sameer and Ayman Shouman Use of Clonidine in Peribulbar Block in Patients Undergoing Cataract Surgery Australian Journal of Basic and Applied Sciences, 5(2): 247-250, 2011 ISSN 1991-8178
- (4) Marta F. G. Barioni, Gabriela R. Lauretti, Argemiro Lauretti-Fo, Newton L. Pereira, B Pharm, Peribulbar vs. oral clonidine in eye surgery Journal of Clinical Anesthesia 14:140 –145, 2002
- (5) Rashmi Madan, Neerja Bharti, Dilip Shende, Sudershan K. Khokhar, and Hira L. Kaul, A Dose Response Study of Clonidine with Local Anesthetic Mixture for Peribulbar Block: A Comparison of Three Doses Anesth Analg 2001;93:1593–7
- (6) Verda Toprak, Ismet Topcu, Baris Toprak, Zeki Kaygisiz, Esin Baser, Cenap Güler: A Comparison of 1% Ropivacaine and 1% Ropivacaine with Clonidine for Retrobulbar Block. The Internet Journal of Anesthesiology. 2006. Volume 11 Number 1.
- (7) Calenda E, Quintyn JC, Brasseur G. Peribulbar anaesthesia using a combination of lidocaine, bupivacaine and clonidine in vitreoretinal surgery Indian J Ophthalmol. 2002 Sep;50(3):205-8.

## conclusion

---

- (8) Zahoor Ahmad Shah, Raksha Kundal, Arun Kumar Gupta, Mohd Sadiq Malla, Faisal Zahoor, Vijay Kundal, Sameena Ashraf Kirmani, Shigufta Qazi Efficacy of analgesic effect of low dose intrathecal clonidine as adjuvant to bupivacaine in urogenital surgeries: study in a Kashmiri population Sri Lankan Journal of Anaesthesiology 2012 Vol 20, No 1
- (9) Masuki S, Dineno FA, Joyner MJ, Eisenach JH Selective alpha<sub>2</sub>-adrenergic properties of dexmedetomidine over clonidine in the human forearm J Appl Physiol. 2005 Aug;99(2):587-92. Epub 2005 Mar 31..
- (10) Butterworth JF, Strichartz GR. The  $\alpha_2$ -adrenergic agonists clonidine and guanfacine produce tonic and phasic block of conduction in rat sciatic nerve fibers. Anesth Analg 1993;76: 295–301.
- (11) De jong RH, Wagman IH. Physiological mechanisms of peripheral nerve block by local anesthetics. Anesthesiol 1963;24:684-727.
- (12) Shiv Kumar Raghuwanshi, Nupur Chakravarty, Dinesh Prasad Asati, Vishal Bankwar Use of Clonidine as an Adjuvant to Infiltration Anaesthesia in Tympanoplasty: A Randomized Double Blind Study Indian Journal of Otolaryngology and Head & Neck Surgery June 2013
- (13) Colin J. L. McCartney, M.B.Ch.B., F.R.C.A., F.R.C.P.C., Edel Duggan, M.D., F.F.A.R.C.S.I., and Emma Apatu, B.A. Should We Add Clonidine to Local Anesthetic for Peripheral Nerve Blockade? A Qualitative Systematic Review of the Literature Regional Anesthesia and Pain Medicine Vol. 32 No. 4 July–August 2007
- (14) Eledjam JJ, Deschodt J, Viel EJ, Lubrano JF, Charavel P, d'Athis F, du Cailar J. Brachial plexus block with bupivacaine: effects of added alpha-adrenergic agonists: comparison between clonidine and epinephrine. Can J Anaesth. 1991 Oct;38(7):870-5.
- (15) Hutschala D, Mascher H, Schmetterer L, Klimscha W, Fleck T, Eichler HG, Tschernko EM (2004) Clonidine added to bupivacaine enhances and prolongs analgesia after brachial plexus block via a local mechanism in healthy volunteers. Eur J Anaesthesiol 21:198-204
- (16) YaDeau JT, Lasala VR, Paroli L, Kahn RL, Jules-Elysee KM, Levine DS, Wukovits BL, Lipnitsky JY: Clonidine and analgesic duration after popliteal fossa nerve blockade: randomized, double-blind, placebo- controlled study. Anesth Analg 2008, 106:1916-1920.

## conclusion

---

- (17) Dobrydnjov I, Axelsson K, Gupta A, Lundin A, Holmstrom B, Granath B. Improved analgesia with clonidine when added to local anesthetic during combined spinal-epidural anesthesia for hip arthroplasty: a double-blind randomized and placebo-controlled study. *Acta Anaesthesiol Scand* 2005; 49:538-45.
- (18) Bajwa SJ, Bajwa SK, Kaur J, Singh A, Singh A, Parmar SS. Prevention of hypotension and prolongation of postoperative analgesia in emergency cesarean sections: A randomized study with intrathecal clonidine. *Int J Crit Illn Inj Sci.* 2012 ;2(2):63-9..