

## Evaluation of antioxidant potential of 124 Egyptian plants with emphasis on the action of *Punica granatum* leaf extract on rats

<sup>1</sup>Moussa, A. M., <sup>1</sup>Emam, A. M., <sup>1\*</sup>Diab, Y. M., <sup>2</sup>Mahmoud, M. E. and <sup>1</sup>Mahmoud, A. S.

<sup>1</sup>Department of Biochemistry, Faculty of Agriculture, Fayoum University, Egypt  
<sup>2</sup>Department of Agriculture and Chemistry, Faculty of Agriculture, Minia University, Egypt

**Abstract:** There is currently an upsurge of interest in phytochemicals as a new source of natural antioxidants to be used in foods and pharmaceutical preparations to replace synthetic antioxidants, which are being restricted due to their potential health risks and toxicity. The chloroform and methanolic leaf extracts of 124 Egyptian plant species belonging to 56 families were investigated and compared for their antioxidant activity by DPPH scavenging assay. Among the 124 plant species tested, 18 exhibit extremely high antiradical activity (> 80% inhibition). The methanolic leaf extract of the promising plant species (18 plants) were further subjected to determine their IC<sub>50</sub> values and total phenolic and flavonoid contents. The IC<sub>50</sub> ranged from 18.68 to 30.97 µg/ml, while total phenolic and flavonoid contents ranged from 162.06±4.55 to 242.26±18.65 mg Tannic acid equivalent (TAE)/g extract and from 32.56±1.52 to 157.96±5.85 mg Rutin equivalent (RE)/g extract, respectively. Correlation coefficient (r) between the DPPH radical scavenging activity (IC<sub>50</sub>) and the total phenolic and flavonoid contents (r=0.63 and 0.51, respectively) suggested that phenolics and flavonoids in the extracts were partly responsible for the antiradical activities. The edibility and safety limits of leaf methanolic extract of *Punica granatum*, which exerted the highest antioxidant activity was assessed by using experimental albino rats. The results indicated that this extract was edible and safe at concentration of 20, 40 and 60 ppm within 9 weeks.

**Keywords:** Natural antioxidants, DPPH, total phenolic content, total flavonoid content, free radical scavenging activity, plant extracts, rats

### Introduction

Free radicals may be defined as any chemical species that are capable of existing with one or more unpaired outer shell electrons. They are extremely reactive and generally highly unstable (Martinez-Cayuela, 1995). They are produced endogenously during cellular metabolism in all forms of aerobic living system, in addition to exogenous sources such as environmental pollutants, drug, radiation, and pathogens (Ansari, 1997). The imbalance between production of reactive oxygen species (ROS) like O<sub>2</sub><sup>-</sup>, H<sub>2</sub>O<sub>2</sub>, OH<sup>-</sup>, ROO<sup>-</sup> and the capacity of the normal detoxification system in favor of the oxidants lead to oxidative stress, which itself lead to cellular damage caused by the interaction of ROS with cellular constituents. Tissue damage resulting from oxidative stress has been implicated in the pathology of a number of disorder diseases such as cancer, inflammatory joint disease, cardiovascular diseases, cataract and could play a role in neurodegenerative diseases and ageing processes (Steer *et al.*, 2002).

The term antioxidant refers to a compound that can delay or inhibit the oxidation of biomolecules by

inhibiting the initiation or propagation of oxidative chain reactions and which can thus prevent damage done to the body's cell by oxygen, i.e. ROS. Epidemiological studies have shown that many phytonutrients particularly phenolic compounds might protect the human body against damage by ROS. The consumption of fruits and vegetables which contain a wide variety of antioxidant phenolic compounds was reported to have potential health benefits (Sumino *et al.*, 2002).

In addition, antioxidants have been widely used in food industry to prolong the shelf life. Synthetic antioxidants such as butylated hydroxy toluene (BHT) and butylated hydroxy anisole (BHA) are widespread food additives used to preserve against deterioration; however their use is increasingly restricted due to their potential health risks and toxicity. Moreover, there is a growing awareness among consumer regarding food additives safety (Valentao *et al.*, 2002). These reasons may explain the interest in examining plant extracts as a source of safe, cheap and effective antioxidants and the growing interest in nutraceuticals.

Plant derived antioxidants exert their effects by enhancing the level of antioxidant enzymes such as

\*Corresponding author.

Email: [ydiab@hotmail.com](mailto:ydiab@hotmail.com)

Tel: +20105310066; Fax: +20 84 633 5153

superoxide dismutase or by lowering the levels of lipid peroxides in the blood or liver (Usoh *et al.*, 2005). It is recognized that antioxidant (mainly phenolic) compounds from plant extracts can act by either free radical scavenging, singlet oxygen quenching, chelating of transitional metal such as iron, as well as a reducing agents and activator of antioxidative defense enzyme systems to suppress radical damage in biological system (Lodovici *et al.*, 2001).

Phenolic compounds are secondary plant metabolites that possess on their structure a benzenic ring substituted by, at least, one hydroxyl group (Manach *et al.*, 2004). These compounds are ubiquitously distributed in the plant kingdom and exhibit a wide range of pharmacological and medicinal properties, including anti-inflammatory, anti-carcinogenic, vasodilatory actions (Middleton *et al.*, 2000); these protective effects have been mostly ascribed to their free radical scavenging, metal chelating and antioxidant action.

In the recent years a large number of endemic plant species have been screened as a viable source of phenolic antioxidants (Wong *et al.*, 2006; Tawaha *et al.*, 2007; Annan and Houghton 2008; Dall'Acqua *et al.*, 2008, Borneo *et al.*, 2009; Rohman *et al.*, 2010).

The aim of this study was to evaluate *in vitro* antioxidant potential of the leaves of 124 plant species growing in Egypt. The effect of three concentrations of methanolic extract which showed the highest antioxidant potential i.e. *Punica granatum* leaf extract on body weight, liver and kidney functions of albino rats was also investigated to assess the safety limits of this extract.

## Materials and Methods

### Chemicals

All chemicals and reagents used in this study were of analytical grade purchased from Sigma Chemical Co. (St.Louis,MO,USA) Aldrich Chemical Co. (Steinehein, Germany), BDH(Dorset-England) or Fluka Chemie Co. (Buchs, Switzerland).

### Plant material

A total of 124 leaf samples of plant species belonging to 56 families were collected during April and May 2008 from El-Shrouk farm, Cario-Alexandria desert road at 72 km north of Cario. The botanical identification of the collected plants was authenticated by Dr. T. Labeb, Herbarium of Orman garden, Horticulture Research Institute, Giza, Egypt. A voucher specimen of each plant was deposited in the Herbarium of the Biochemistry Department,

Faculty of Agriculture, Fayoum University. The leaves were cleaned, air-dried in the shade, and then powdered by laboratory mill to 24 mesh. Powdered materials were maintained in an air tight container at room temperature ( $28\pm 2^{\circ}\text{C}$ ), and protected from light until use.

### Extraction

A known amount of air dried powdered leaves of each plant sample was extracted at room temperature ( $28\pm 2^{\circ}\text{C}$ ) with chloroform then with methanol. This procedure was repeated at least five times until each organic solvent became colorless. The obtained extracts were filtered over Whatman No.1 filter paper and the combined extract (filtrate) was evaporated to dryness under vacuum at  $45^{\circ}\text{C}$  using a rotary evaporator. The dried extract (residue) obtained from each plant material was stored in a desiccators at  $4^{\circ}\text{C}$  until further use.

### DPPH free radical scavenging assay

Free radical scavenging activity was measured by the stable 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical, according to the method described by Brand-Williams *et al.*, (1995). Briefly, a 2.0 ml of methanolic solution of each extract at concentration of  $50\ \mu\text{g/ml}$  was added to a 2.0 ml solution of DPPH (25 mg/l) in methanol, and the reaction mixture was shaken vigorously. After incubation at room temperature ( $28\pm 2^{\circ}\text{C}$ ) for 30 min, the absorbance (A) of DPPH for control and samples was determined by Cecil 3000 spectrophotometer at 517nm, and the radical scavenging activity of each extract was expressed as percentage inhibition of free radical DPPH.

$$\% \text{ inhibition (\% Anti-radical activity)} = [(A \text{ control} - A \text{ sample}) / A \text{ control}] \times 100$$

Triplicate measurements were taken and mean values calculated.  $\text{IC}_{50}$  (concentration of each sample at 50% inhibition) was obtained by linear regression analysis of the dose response curve, plotted between % inhibition and concentrations ( $2.5\text{-}100\ \mu\text{g/ml}$ ) L-Ascorbic acid was used as a positive control.

### Determination of total phenolic content

The total phenolic content was determined using the Folin-Ciocalteu reagent as described by Yu *et al.* (2002). Two hundred micro liters of each sample were mixed with  $500\ \mu\text{l}$  of the Folin-Ciocalteu reagent and 1.5 ml of 20% sodium carbonate. The mixture was shaken thoroughly and made up to 10 ml using distilled water. After 2 h of incubation at room temperature, the absorbance of the resulting blue-colored solution was measured at 765 nm. The assay was carried out in triplicate and the mean values were calculated.

The concentration was determined from the standard curve prepared using serial concentrations of standard tannic acid solution. Total phenolic contents in plant extracts were expressed as mg tannic acid equivalent (TAE)/g plant extract.

#### *Determination of total flavonoid content*

Total flavonoid content was determined using the method described by Lamaison and Carnet, 1990 as follows: Two hundred micro liters of each sample were transferred to a test tube and evaporated to dryness. To the residue, 5 ml of 0.1 M AlCl<sub>3</sub> were added and shaken. The intensity of the developed color was measured at 415 nm after 40 min versus the prepared blank. Triplicate measurements were taken and mean values calculated. Rutin was used as standard for the calibration curve. Total flavonoid content was expressed as mg rutin equivalent (RE) /g plant extract.

#### *Effect of the methanolic extract of Punica granatum on albino rats*

The effect of orally administrating of *Punica granatum* methanolic leaf extract on body weight, liver, and kidney function tests of albino rats was studied as follows:

#### *Animals and diets*

Twenty female albino rats with body weight of 95-101 g were obtained from the Faculty of Agriculture, Menia University, Egypt, and were acclimated for one week prior to the experiment. They were housed in groups of five each in universal polypropylene cages at room temperature (25±2°C) and at a photoperiod 12 h/day. Animals were fed on standard laboratory chow (El-Nasar Lab. Chem. Co., Egypt) and water *ad libitum* till the end of the experiment (9 weeks).

#### *Experimental design*

The rats were divided into four experimental groups. Group I rats (control group) orally received 1 ml H<sub>2</sub>O/day. Group II rats orally received daily 20 ppm methanolic *Punica granatum* extract (in 1 ml H<sub>2</sub>O). Group III rats orally received daily 40 ppm methanolic *Punica granatum* extract (in 1 ml H<sub>2</sub>O). Group IV rats orally received daily 60 ppm methanolic *Punica granatum* extract (in 1 ml H<sub>2</sub>O). Body weight (BW) was recorded weekly during the experimental period (9 weeks) and food intake was measured daily during the adaptation and the experimental periods.

At the end of the experimental period (9 weeks) animals were fasted overnight, blood samples were withdrawn by a fine capillary glass tubes from the orbital plexus vein. The blood was collected in EDTA

containing tubes and left at room temperature. Plasma was obtained by centrifugation at 3000 rpm for 15min and then stored at -20°C until analysis. Liver and kidney organs were excised, rinsed in chilled saline solution, and then plotted on filter paper, weighed separately to calculate the absolute organs weight.

#### *Analytical procedures*

Plasma enzyme activity of aspartate aminotransferase (AST; E.C.2.6.1.1.) and alanine aminotransferase (ALT; E.C.2.6.1.2.) activities were measured according to the method described by Reitman and Frankel, 1957. Total protein, albumin, bilirubin and urea were determined according to the methods described previously (Gornal *et al.*, 1948; Doumas *et al.*, 1971; Fawcett and Soctt, 1960), using enzymatic colorimetric procedures Kits from Bio-Diagnostic Co., Egypt. Globulin was determined by difference between total protein and albumin.

#### *Statistical analysis*

Statistical Package for Social Science (SPSS) window's version (10, USA) was used for analysis of the data.

## **Results and Discussion**

#### *DPPH radical scavenging activity*

The chloroform and methanol leaf extracts of 124 plant species extracts at 50 µg/ml were tested for antioxidant activity using the DPPH assay. The results were expressed as percentage inhibition (Table 1). As shown in Table 1, there is a wide range of free radical scavenging activity of the chloroform and methanolic extracts of the plant species analyzed. The values ranged from 0.5 to 49% and from 3 to 96% for chloroform and methanolic extracts, respectively. The variation of the free radical scavenging activity may be due to the differences in their secondary constituents (Sudjaroen *et al.*, 2005).

According to the data obtained the plant species were classified into four categories on the basis of free radical scavenging activity low antioxidant power (<25% inhibition), medium (25-50% inhibition), high (50-80% inhibition) and extremely high (>80% inhibition). The number of plants in each of the low, medium, high and extremely high antioxidant power categories was 44, 52, 11 and 18 respectively.

The extremely high antioxidant power category (18 plant species) was further subjected to determine their IC<sub>50</sub> values, which is defined as the concentration of substrate at 50% inhibition (Han *et al.*, 2004). The IC<sub>50</sub> values (µg/ml) of DPPH radical scavenging

**Table 1.** 124 Egyptian plants studied and free radical scavenging activity of their chloroform and methanolic leaf extracts

Plant Species	Family	% Antiradical activity Chloroform extract (50 µg/ml)	Methanol extract (50 µg/ml)
<i>Adhatoda vasica</i>	Acathaceae	3	13
<i>Dracaena marginata</i>	Agavaceae	15	13
<i>Yucca desmetiana</i>	"	3	6
<i>Polygonum tuberosum</i>	"	8	12
<i>Alternanthera versicolor</i>	Amaranthaceae	0.5	71
<i>Amaranthus tricolor</i>	"	1	0.0
<i>Crinum longifolium</i>	Amaryllidaceae	6	21
<i>Schinus molle</i>	Anacardiaceae	15	12
<i>Foeniculum vulgare</i>	Apiaceae	46	16
<i>Carissa grandiflora</i>	Apocynaceae	2	0.0
<i>Thevetia peruviana</i>	"	26	29
<i>Plumeria alba</i>	"	28	15
<i>Tabernaemontana divaricata</i>	"	40	44
<i>Vinca rosea</i>	"	9	48
<i>Nerium oleander</i>	"	19	31
<i>Anthurium scherzerianum</i>	Araceae	14	37
<i>Sciadophyllum pulchrum</i>	Araliaceae	1.5	18
<i>Schefflera actinophylla</i>	"	3	94
<i>Schefflera arboricola</i>	"	2	0.0
<i>Cryptostegia grandiflora</i>	Asclpiadaceae	0.0	35
<i>Chrysanthemum Red</i>	Asteraceae	15.5	35
<i>Santolina chamaecyparissus</i>	"	11	46
<i>Leucaena capensis</i>	Bignoniaceae	0.0	55
<i>Spathodea nilotica</i>	"	19	85
<i>Kigelia pinnata</i>	"	3	0.0
<i>Jacaranda acutifolia</i>	"	18	55
<i>Leucaena capensis</i>	"	9	90
<i>Bombax malabaricum</i>	Bombacaceae	2	96
<i>Cordia sebestena</i>	Boraginaceae	15	46
<i>Poinciana regia</i>	Caesalpiniaceae	15	15
<i>Lonicera japonica</i>	Caprifoliaceae	5	16
<i>Euonymus japonica</i>	Celastraceae	3	66
<i>Conocarpus erectus</i>	Combretaceae	15	45
<i>Terminalia arjuna</i>	"	27	42
<i>Tradescantia zebrina</i>	Commelinaceae	1	16
<i>Tradescantia spp</i>	"	44	0.0
<i>Gazania splendens</i>	Compositae	0.0	89
<i>Chrysanthemum frutescens</i>	"	0.0	87
<i>Aspidistra lurida</i>	Convallariaceae	0.0	29
<i>Thuja orientalis</i>	Cupressaceae	0.0	71
<i>Cupressus sempervirens</i>	"	6	65
<i>Cupressus macrocarpa</i>	"	3	48
<i>Cyperus alternifolius</i>	Cyperaceae	2	31
<i>Elaeagnus macrophylla</i>	Elaeagnaceae	45	8
<i>Jatropha integerrima</i>	Euphorbiaceae	13	19
<i>Jatropha multifida</i>	"	47	17
<i>Euphorbia splendens</i>	"	25	31
<i>Acalypha marginata</i>	"	29	80
<i>Sesbania aegyptiaca</i>	Fabaceae	17	43
<i>Pithecellobium dulce</i>	"	11	23
<i>Pelargonium zonal</i>	Geraniaceae	12	32
<i>Pelargonium odoratissimum</i>	"	9	89
<i>Rosmarinus officinalis</i>	Lamiaceae	28.6	31
<i>Salvia officinalis</i>	"	45	49
<i>Lavandula angustifolia</i>	"	26	36
<i>Laurus nobilis</i>	Lauraceae	49	89
<i>Cordyline fruticosa</i>	Loganiaceae	18	91
<i>Bauhinia variegata</i>	Leguminosae	18	94
<i>Cassia didymobotrya</i>	"	16.5	12
<i>Cassia fistula</i>	"	21.6	45
<i>Sansevieria guineensis</i>	Liliaceae	23.5	18
<i>Ruscus hypoglossum</i>	"	0.0	42
<i>Phalangium variegata</i>	"	2	17
<i>Asparagus setaceus</i>	"	2	15
<i>Asparagus plumosus</i>	"	0.0	16
<i>Buddleia hybrida</i>	Loganiaceae	10	50
<i>Caesalpinia pulcherrima</i>	"	25	27
<i>Coratonia siliqua</i>	"	15	40
<i>Albizia anthelmintica</i>	"	20	49
<i>Parkinsonia aculeata</i>	"	17	9
<i>Lagerstroemia indica</i>	Lythraceae	28	89
<i>Hibiscus mutabilis</i>	Malvaceae	0.0	40
<i>Althaea rosea</i>	"	19	58
<i>Khaya senegalensis</i>	Meliaceae	13	86
<i>Melia azadirach</i>	"	3	32
<i>Calliandra haematocephala</i>	Mimosaceae	10	23
<i>Albizia stipulata</i>	"	15	18
<i>Dracaena fragrans</i>	Moraceae	1	33
<i>Morus alba</i>	"	14	40
<i>Ficus nitida</i>	"	5	25
<i>Ficus benjamina</i>	"	2	37
<i>Ficus elastica Var decora</i>	"	6	88
<i>Ficus hawaii</i>	"	3	35
<i>Ficus ali</i>	"	1	3
<i>Ficus relogiosa</i>	"	3	24

Plant Species	Family	% Antiradical activity Chloroform extract (50 µg/ml)	Methanol extract (50 µg/ml)
<i>Ficus elastica</i>	"	11.5	41
<i>Ficus enfiectoria</i>	"	3.5	31
<i>Ficus Natalia</i>	"	5	16
<i>Moringa pterygosperma</i>	Moringaceae	15	30
<i>Myoporum pictum</i>	Myoporaceae	42.7	26
<i>Eucalyptus rostrata</i>	Myrtaceae	9	90
<i>Eugenia uniflora</i>	"	13	16
<i>Myrtus Communis</i>	"	26.6	90
<i>Calistemon lanceolatus</i>	"	23	78
<i>Bougainvillea spectabilis</i>	Nyctaginaceae	12	38
<i>Bougainvillea "pixie-pink"*</i>	"	22	39
<i>Bougainvillea glabra var. sanderjana a</i>	"	15.6	0.0
<i>Bougainvillea glabra "variegata"</i>	"	20	50
<i>Narcissus tazetta</i>	Oleaceae	22	40
<i>Jasminum primulinum</i>	"	6	30
<i>Jasminum grandiflorum</i>	"	39	52
<i>Ligustrum ovalifolium</i>	"	9	11
<i>Jasminum sp</i>	"	45.7	34
<i>Araucaria excelsa</i>	Pinaceae	22	43
<i>Pittosporum tobira var. variegata</i>	Pittosporaceae	18.6	54
<i>Pittosporum tobira</i>	"	11	9
<i>Grass sp</i>	Poaceae	5	30
<i>Nephrleps bostoniensis</i>	Polypodiaceae	16	43
<i>Punica granatum</i>	Punicaceae	9	96
<i>Rosa spp</i>	Rosaceae	26.6	27
<i>Rosa banksiae</i>	"	25	42
<i>Didonia viscosa</i>	Sapindaceae	0.0	90
<i>Hydrangea Red</i>	Saxifragaceae	4	29
<i>Leucophyllum frutescens</i>	Scrophulariaceae	16.4	31
<i>Russelia juncea</i>	"	17	30
<i>Datura arborea</i>	Solanaceae	17	56
<i>Cestrum diurnum</i>	"	30	50
<i>Sterculia diversifolia</i>	Sterculiaceae	14	88
<i>Brachichton acerifolium</i>	"	16	22
<i>Strelitzia reginae</i>	Streliziaceae	32	45
<i>Duranta repens</i>	Verbenaceae	16	44
<i>Lantana camara</i>	"	12	32
<i>Lantana montevidensis</i>	"	11	30
<i>Vitex trifolia "purpurea"</i>	"	22	89

activity from dried methanolic extracts of these plant species and the positive control (Ascorbic acid) are presented in Table 2. As shown in this Table, *Punica granatum* and *Bombax malabaricum* showed the highest DPPH radical scavenging activity (the lowest values of  $IC_{50}$  i.e., 18.68 and 18.80 µg/ml, respectively) while the plant with lowest DPPH scavenging activity (the highest value of  $IC_{50}$  i.e., 30.97) was *Spathodea tilotica*.

Between those high and low values a narrow range of antioxidant abilities were obtained among the 18 tested plant species. The DPPH radical scavenging effect of the positive control i.e. ascorbic acid was higher ( $IC_{50}$  = 11.5 µg/ml) than all the 18 plant species studied. The  $IC_{50}$  values of the analyzed plant species were roughly from 1.62 to 2.69 times lower than ascorbic acid. However we should keep in mind that those of high DPPH scavenging activity value obtained for ascorbic acid is because we performed the assay on highly purified reference standard and not in complex materials such as that of plant species analyzed. This implies that although there are plants with good antioxidant abilities further concentration or purification is needed to achieve better antioxidant capacities.

In the recent years various plant species have been tested as botanical antioxidants using DPPH assay (Wong *et al.*, 2006; Annan and Houghton 2008; Dall'Acqua *et al.*, 2008; Borneo *et al.*, 2009; Rohman

*et al.*, 2010). These species exhibited a broad range of antioxidant activities. The highest antioxidant activity has been reported for *Rupus ulmifolius*,  $IC_{50}$  values 5.1 µg/ml (Dall'Acqua *et al.*, 2008), whereas the lowest antioxidant was reported for *Thelesperma megapotamicam*,  $IC_{50}$  values 2000 µg/ml (Borneo *et al.*, 2009).

#### Total phenolics and flavonoids contents

The systematic literature collection, pertaining to this investigation indicates that the plant phenolics constitute one of the major groups of compounds acting as primary antioxidant or free radical terminators. Therefore, it is worthwhile to determine their total amount in plants chosen for the study. Flavonoids one of the most diverse and widespread group of natural compounds, are likely to be the most important natural phenolics due to their broad spectrum of chemical and biological activities, including antioxidant and free radical scavenging properties (Kahkonen *et al.*, 1999). Therefore the contents of flavonoids is also determined.

Total phenolics content of methanolic extracts (TPC) of the 18 promising plant species were expressed as tannic acid equivalent (TAE) in milligrams per gram extract. As shown in Table 2, the TPC ranged from 162.06 to 242.26 mg TAE/g extract. *Punica granatum* and *Myrtus communis* showed the highest TPC of 242.26 and 241.56 mg TAE/g extract,

while *Vitex trifolia* "purpurea" was the lowest one in TPC (162.02 mg TAE/g extract).

The total flavonoid content (TFC) of the methanolic extracts of the promising plant species was expressed as milligram rutin equivalents (RE) per gram extract and are presented in Table 2.

**Table 2.** Total phenolic and total flavonoid contents of leaf methanolic extracts of the promising antioxidant plant species

Total flavonoids (mg RE/g extract)	Total Phenolic (mg TAE/g extract)	IC <sub>50</sub> (µg/ml)	Family	Plant Scientific name
54.360 ±5.070 <sup>m</sup> 157.96	242.26 ±18.65 <sup>a</sup> 214.06	18.68	Punicaceae	<i>Punica granatum</i>
±5.850 <sup>n</sup> 32.560	±5.050 <sup>b,c</sup> 220.86	18.80	Bombacaceae	<i>Bombax malabaricum</i>
±1.520 <sup>n</sup> 109.76	±4.750 <sup>ab</sup> 226.00	19.23	Araliaceae	<i>Schefflera actinophylla</i>
±0.850 <sup>h</sup> 67.760	±7.140 <sup>ab</sup> 220.26	19.36	Leguminosae	<i>Bauhenia variegata</i>
±9.500 <sup>l</sup> 113.66	±33.85 <sup>ab</sup> 208.26	20.73	Myrtaceae	<i>Eucalyptus rostrata</i>
±1.050 <sup>e</sup> 126.96	±41.65 <sup>abcd</sup> 241.56	21.05	Sapindaceae	<i>Didonia viscosa</i>
±1.150 <sup>d</sup> 111.06	±11.75 <sup>a</sup> 226.40	21.53	Myrtaceae	<i>Myrtus Communis</i>
±5.500 <sup>ef</sup> 135.66	±23.07 <sup>ab</sup> 162.06	22.55	Bignoniaceae	<i>Tecomaria capensis</i>
±0.550 <sup>e</sup> 48.06	±4.550 <sup>f</sup> 208.86	22.94	Verbenaceae	<i>Vitex trifolia</i> "purpurea"
±0.350 <sup>n</sup> 107.06	±10.95 <sup>abcd</sup> 223.16	23.57	Compositae	<i>Gazania splendens</i>
±0.450 <sup>h</sup> 71.36	±10.85 <sup>ab</sup> 177.16	74	Lythraceae	<i>Lagerstroemia indica</i>
±0.650 <sup>k</sup> 88.76	±6.250 <sup>def</sup> 205.16	24.57	Euphorbiaceae	<i>Acalypha marginata</i>
±0.850 <sup>l</sup> 104.96	±9.150 <sup>abcde</sup> 180.66	25.30	Lauraceae	<i>Laurus nobilis</i>
±0.550 <sup>h</sup> 111.06	±12.45 <sup>def</sup> 171.66	26.00	Geraniaceae	<i>Pelargonium oderatissimum</i>
±0.550 <sup>ef</sup> 69.06	±2.450 <sup>ef</sup> 208.66	27.32	Sterculiaceae	<i>Sterculia diversifolia</i>
±0.250 <sup>kl</sup> 149.36	±8.650 <sup>abcd</sup> 215.06	27.83	Compositae	<i>Chrysanthemum frutescence</i>
±0.550 <sup>h</sup> 129.96	±43.05 <sup>abc</sup> 197.26	29.61	Meliaceae	<i>Khaya senegalensis</i>
±0.750 <sup>d</sup>	±6.550 <sup>bcd</sup>	30.97	Bignoniaceae	<i>Spathodea tilotica</i>
<b>11.5</b>				<i>Ascorbic acid</i>

Each value is presented as mean ±SD (n=3). Means within each column with different letters (a-m) differ significantly (p<0.05).



The TFC ranged from 32.56 to 157.96 mg RE/g extract for *Bombax malabaricum* and *Schefflera actinophylla*. The 18 analyzed plant species have the same geographic origin and grow in the same natural condition; nevertheless the plants belonging to different families. It is well known that the amount of total phenolics vary with respect to families and varieties (Kahkonen *et al.*, 1999; Djeridane *et al.*, 2006).

The results in Table 2, also indicated that *Punica granatum* had the best DPPH radical scavenging activity and also the highest total phenolics content. However, as previously mentioned, *Myrtus communis* had higher phenolics content (241.56 mg TAE/g extract) than *Bombax malabaricum*, *Schefflera actinophylla*, *Bauhenia variegata*, *Eucalyptus rostrata* and *Didonia viscosa* (214.06, 220.86, 226.0, 220.26 and 208.26 mg TAE/g extract, respectively), but these plant extracts exhibited high DPPH radical scavenging activity than *Myrtus communis*. Also *Vitex trifolia* had lower phenolics content (162.06 mg TAE/g extract) than *G.splendens*, *L.indica*, *A. marginata*, *L. nobilis*, *P. oderatissimum*, *S.diversifolia*, *C. frutescence*, *K. senegalensis* and *S. titotica* (208.86,

223.16, 177.16, 205.16, 180.66, 171.66, 208.66, 215.06 and 197.26 mg TAE/g extract respectively, however it had high DPPH radical scavenging activity. Folin assay gives a crude estimate of the amount of phenolic compounds present in an extract. It is not specific to polyphenols but many interfering compounds may react with the reagent, giving elevated apparent phenolic concentrations (Singleton *et al.*, 1999). Moreover, various phenolics compounds respond differently in this assay, depending on the number of phenolic groups they have and total phenolics content does not incorporate necessarily all the antioxidant that may be present in an extract (Tawaha *et al.*, 2007).

#### Correlation between DPPH radical scavenging activity and total phenolic and flavonoid compounds

The correlation coefficient (r) between the free radical scavenging activity and both the total phenolic and the total flavonoid contents of the promising plants was determined (Figure 1). The correlation coefficient between the scavenging activity obtained from DPPH assay and the phenolic content was 0.63 (Figure 1A) whereas with total flavonoid contents has correlation coefficient of 0.51 (Figure 1B). These results suggested that phenolic and flavonoid compounds contributed of 63.0% and 51.0% to free DPPH radical scavenging of the analyzed plant extracts. Also, it can be stated that scavenging effects is not limited to phenolic and flavonoid compounds. The activity also comes from the presence of other antioxidant secondary metabolites in the extracts which directly or indirectly contribute to the activity.



**Figure 1.** Correlation between free DPPH radical scavenging activity (IC<sub>50</sub>) and total phenolic contents (A) and with total flavonoid contents (B)

**Table 3.** Effect of different levels of orally methanolic extract administration of *Punica granatum* on body weight and liver and kidney organs weight

Group No.	Initial body Weight (g)	Final body Weight (g)	Liver Weight (g)	Kidney Weight (g)	Weight gain (g)	Food consumption (g/day)	Food efficiency
I (control)	100.8±12.7	192.0±11.27	6.23±0.41	1.23±0.14	91.2±11.27	82.10	1.11
II (20mg/kg)	95.0±0.0	195.67±0.58	7.09±0.26	1.34±0.22	100.7±0.58	86.30	1.17
III (40mg/kg)	99.0±2.24	196.0±1.73	6.58±0.5	1.31±0.07	97.00±2.54	85.04	1.14
IV (60mg/kg)	99.0±2.24	196.0±1.73	7.02±0.41	1.28±0.09	97.00±2.54	91.36	1.06

Values are expressed as mean ± S.D. (n=5).

**Table 4.** Effect of different levels of orally methanolic extract administration of *Punica granatum* on liver and kidney functions in plasma of female albino rats

Group No.	ALT(GPT) (U/L)	AST(GOT) (U/L)	T.Protein (g/dl)	Albumin (g/dl)	Globulin (g/dl)	Urea (mg/dl)	Total bilirubin (mg/dl)
I (control)	42.9±2.7	62.7±7.2	5.902±0.62	3.689±0.20	2.213±0.67	79.076±24.69	0.567±0.16
II (20mg/kg)	42.6±3.2	61.38±6.2	5.408±0.54	3.775±0.84	1.633±0.79	78.743±23.17	0.308±0.03
III (40mg/kg)	44.1±4.6	59.4±9.6	5.845±0.74	4.087±0.64	1.758±0.97	64.831±15.24	0.518±0.31
IV (60mg/kg)	42.6±3.4	62.92±7.3	5.55±0.50	3.90±0.57	1.654±0.81	65.423±21.75	0.254±0.10

Values are expressed as mean ± S.D. (n=5).

These data are in accordance others (Ou *et al.*, 2003; Rohman *et al.*, 2010; Cox *et al.*, 2010) who have reported that there is no correlation between the content of these main antioxidant compounds and the radical scavenging activity, whereas many studies indicate a linear relationship between these two variables (Djeridane *et al.*, 2006; Tawaha *et al.*, 2007; Borneo *et al.*, 2009).

#### Effect of leaf methanolic extract of *Punica granatum* on albino rats

In order to assess the edibility and safety limits of leaf methanolic extract of *Punica granatum* that, exerted the highest antioxidant activity. Three levels of methanolic extract preparation, including 20, 40 and 60 mg / kg body weight (group II, III and IV respectively) were used in this experiment in addition to control (group I). The obtained results are shown in Table (3) and (4). No significant differences were noted in the mean value of body weight and organs weight among different experimental groups (Table 3). Results obtained in Table 4 showed no significant changes in liver enzymes activity ALT or AST in treated groups compared with control rats. Also, no significant effects were observed in plasma bilirubin, total protein, albumin and globulin in treated groups corresponding to control group. Meanwhile, slight decrease in urea was shown in the different treated groups (Table 4), but this slight decrease is not significant as compared with control group. These

results indicated that orally administration of the three concentrations of leaf methanolic extract of *Punica granatum* has no negative effect on body weight or liver and kidney function tests of female albino rats. The edibility of the methanolic extract was observed as the food consumption and so body weight was increased. The results of the experimental animals indicated that the three concentrations of leaf methanolic extract of *Punica granatum* was safe and edible.

#### Conclusion

In conclusion, this study suggests that some of the plant species studied may be a good source of natural antioxidants. Further studies are warranted for isolation and characterization of the active components of promising plant species as well as *in vivo* evaluation of toxicity of these active components in animals studies.

#### References

- Anasari, K. N. 1997. The free radicals -the hidden culprits- an update. Indian Journal of Medical Sciences 51: 319-336.
- Annan, K. and Houghton, P. 2008. Antibacterial, antioxidant and fibroblast growth stimulation of aqueous extract of *Ficus asperifolia* Miq. and *Gossypium arboreum* L., Wound-healing plants of Ghana. Journal of

- Ethnopharmacology 119: 141-144.
- Borneo, R., Leon, A., Aguirre, A., Ribotta, P. and Cantero, J. 2009. Antioxidant capacity of medicinal plants from the province of Cordoba (Argentina) and their *in-vitro* testing in a model food system. Food Chemistry 112: 664-670.
- Brand-Williams, W.; Cuvelier, M. E. and Berset, C. 1995. Use of a free radical method to evaluate antioxidant activity. Lebensmittel Wissenschaft und Technologie 28: 25-30.
- Cox, S., Abu-Ghannam, N. and Gupta, S. 2010. An assessment of the antioxidant and antimicrobial activity of six species of edible Irish seaweeds. International Food Research Journal 17: 205-220.
- Dall'Acqua S.; Cerrellati, R.; Loi, M. C. and Innocenti, G. 2008. Evaluation of *in-vitro* antioxidant properties of some traditional Sardinian medicinal plants: Investigation of the high antioxidant capacity of *Rubus ulmifolius*. Food Chemistry 106: 745-749.
- Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P. and Vidal, N. 2006. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. Food Chemistry 97: 654-660.
- Doumas, B. T., Watson, W. A. and Biggs, H. G. 1971. Albumin standards and the measurement of serum albumin with bromocresol green. Clinica Chimica Acta 31: 87-96.
- Fawcett, J. K. and Soctt, J. E. 1960. Enzymatic, colorimetric method for determination urea in serum, plasma and urine. Journal of Clinical Pathology 13: 156-162
- Gornall, A. G., Bardawill, C. J. and David, M. M. 1948. Determination of serum protein by means of the biuret reaction. Journal of Biological Chemistry 177: 751-766.
- Han, S. S., Lo, S. C.; Choi, Y. W., Kim, J. H. and Baek, S. H. 2004. Antioxidant activity of crude extract and pure compounds of *Acer ginnala* Max. Bulletin of the Korean Chemical Society 25(3): 389-391.
- Kahkonen, M. P., Hopia, A. I., Vuorela, H. J., Rauha, J. P., Pihlaja, Kujala, T. S. 1999. Antioxidant activity of plant extracts containing phenolic compounds. Journal of Agricultural and Food Chemistry 47: 3954-3962.
- Lamaison, J. L. C. and Carnet, A. 1990. Teneurs en principaux flavonoides des fleurs de *Crataegus monogyna* Jacq et de *Crataegus Laevigata* (Poirlet D.C) en fonction de la vegetation pharmaceut. Acta Helve 65: 315-320.
- Lodovici, M., Guglielmi, F., Casalini, C., Meoni, M., Cheyiner, V. and Dolara, P. 2001. Antioxidant and radical scavenging properties in vitro of polyphenolic extracts from red wine. European Journal of Nutrition 40: 74-77.
- Manach, C., Scalbert, A. Morand, C., Rémés, C. and Jimenez, L. 2004. Polyphenols: Food sources and bioavailability. American Journal of Clinical Nutrition 79: 727-747.
- Martinez-Cayuela, M. 1995. Review: oxygen free radicals and human disease Biochimie 77: 147-147.
- Middleton, Jr. E., Kandaswami, C. and Theoharides, T. C. 2000. The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease and cancer. Pharmacological Reviews 52:673-839.
- Ou, B., Huang, D., Hampsch-Woodili, M. and Flanagan, J. A. 2003. When the east meets west: The relation Yin-Yang and antioxidation-oxidation. Journal of the Federation of American Societies for Experimental Biology FASEB 17: 127-129.
- Reitman, S. and Frankel, S. 1957. A colorimetric method for the determination of serum glutamic-oxaloacetic and glutamic pyruvic transaminases. American Journal of Clinical Pathology 28: 56-63.
- Rohman, A., Riyanto, S., Yuniarti, N., Saputra, W. R., Utami, R. and Mulatsih, W. 2010. Antioxidant activity, total phenolic and total flavonoids of extracts and fractions of red fruit (*Pandanus conoideulam*). International Food Research Journal 17: 97-106.
- Singelton, V. L., Orthofer, R. and Lamuea-Raventos, R. M. 1999. Analysis of total phenolic and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods in Enzymology 29: 152-178.
- Steer, P., Millgard, J., Sarabi, D., Wessby, B. and Kahan, T. 2002. Cardiac and vascular structure and function are related to lipid peroxidation and metabolism. Lipids 37: 231-236.
- Sudjaroen, Y., Haubner, R., Wartele, G., Hull, W. E., Erben, G., Spiegelhalder, B., Changbumrung, S., Bartsch, H. and Owen, R.W. 2005. Isolation and structure elucidation of phenolic antioxidants from Tamarind (*Tamarindus indica* L.) seeds and pericarp. Food and Chemical Toxicology 43: 1673-1682.
- Sumino, M., Sekin, T., Ruanagrunsi, N., Igarashi, K. and Ikkegami, F. 2002. Ardisiphenols and other antioxidant principles from the fruits of *Ardisia colorata*. Chemical and Pharmaceutical Bulletin 50: 1484-1487.
- Tawaha, K., Alali, F. Q., Gharaibeh, M., Mohammad, M. and El-Elimat, T. 2007. Antioxidant activity and total phenolic content selected Jordanian plant species. Food Chemistry 104: 1378-1378.
- Usoh, I., Akpan, E., Etim, E. and Farombi, E. 2005. Antioxidant action of dried flower extracts *Hibiscus sabdariffa* L. on sodium arsenite- induced oxidative stress rats. Pakistan Journal of Nutrition 4:135-141.
- Valentao, P., Fernandes, E., Carvalho, F., Andrade, P. B., Seabra, R. M. and Bastos, M. 2002. Antioxidative properties of carbon *Cynara cardunculus* L. infusion against superoxide radical hydroxy radical and hypochlorous acid. Journal of Agricultural and Food Chemistry 50: 4989-4993.
- Wong, S. O., Leong, L. P. and Koh, J. H. W. 2006. Antioxidant activities of aqueous extracts of selected plants. Food Chemistry 99: 775-783.
- Yu, L., Haley, S., Perret, J., Harris, M., Wilson, J. and Qian, M. 2002. Free radical scavenging properties of wheat extracts. Journal of Agricultural and Food Chemistry 50: 1619-1624.