



**Characteristics, and predictive factors of disease severity in
hospitalized patients with SARS-COV-2 in Fayoum governorate,
Egypt: a multicenter study**

**Doaa Y. Ali¹; Marwa A. Ali²; Abdelrahman M. Ahmed³; Wafaa Y. Abdel-
Wahed⁴**

1. Department of Clinical and Chemical Pathology, Faculty of Medicine, Fayoum University, Egypt.
2. Department of Medical Biochemistry and Molecular Biology, Faculty of Medicine, Fayoum University, Egypt
3. Department of Chest Disease and Tuberculosis, Faculty of Medicine, Fayoum Chest Hospital, Al Fayoum, Egypt.
4. Department of Public Health and Community Medicine. Faculty of Medicine, Fayoum University, Al Fayoum, Egypt

Journal Name : Microbes and infectious diseases , Local

Abstract:

Background: Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) pandemic is a serious health problem all over the world including Egypt, thus realizing the predictive factors and disease's characteristics is an essential issue. **Objectives:** To evaluate the characteristics of laboratory-confirmed cases of SARS-CoV-2 infection in Fayoum governorate, Egypt, and to determine the predictive factors of disease severity. **Methods:** One hundred-fourty patients confirmed with SARS-CoV-2 from the Fayoum governorate, Egypt, were collected in this descriptive multicenter study. The subtype classification of SARS-CoV-2 was according to the World Health Organization (WHO) guideline SARS-CoV-2 disease severity classification. Patients were divided into a asymptomatic/non-severe cases group and a severe/critical case group. Each patient was subjected to chest computed tomography (CT), clinical, and laboratory assessment in form of complete blood count, neutrophil-to-lymphocyte ratio (NLR), lymphocyte-to-monocyte ratio (LMR), platelet-to-lymphocyte ratio (PLR), liver function tests, urea, creatinine, C-reactive protein (CRP), serum ferritin, and D-dimer. **Results:** Severe/critical patients were older (52.0 ± 12.6) with a statistical significantly higher rate of diabetes mellitus, hypertension, and tuberculosis (TB) ($p < 0.001$) than non-severe cases. Dyspnea was the most prevalent significant symptom among severe /critical group (87.5%, $p < 0.001$). A negative correlation between radiographic score and oxygenation index ($r = -0.302$, $p = 0.007$). Using the ROC analysis, the area under curve (AUC) was highest for a radiographic score, D-dimer, CRP, ferritin, and NLR, with $p < 0.05$ in severe/critical cases. **Conclusions:** Elevated CRP, D-dimer, serum ferritin, radiograph score, and NLR may contribute to the judgment of SARS-CoV-2 severity, and help



clinicians to evaluate the patient's condition. Co-infection of SARS-CoV-2 and TB can occur, and may progress towards severe SARS-CoV-2

Keywords: SARS-CoV2, Egypt, Tuberculosis, Severe/critical, Predictive factors