

Diagnostic accuracy of global longitudinal strain in prediction of severity and extent of coronary artery stenosis in patients with acute coronary syndrome

مكان وتاريخ النشر:

ACTA CARDIOLOGICA

<https://doi.org/10.1080/00015385.2022.2066810>

ISSN: 0001-5385, 1784973x

ملاحظات

- بحث مستخرج من رسالة
- هذا البحث جارٍ تقييمه حالياً ومقدم للجنة الترقيات لطب الحالات الحرجة من دكتور تامر محمد سيد مدرس الحالات الحرجة
- أسماء الباحثين:

د/ أسامة محمود ممتاز
أستاذ مساعد الحالات الحرجة- كلية الطب - جامعة الفيوم

د/ تامر محمد سيد
مدرس الحالات الحرجة- كلية الطب - جامعة الفيوم
أ.د خالد أحمد الخشاب

أستاذ أمراض القلب - كلية الطب - جامعة الفيوم
د. ايمان محمود عبد الفتاح

مدرس أمراض القلب - كلية الطب - جامعة الفيوم

د/ وسام صلاح عبد الونيس
مدرس الحالات الحرجة- كلية الطب - جامعة الفيوم

ABSTRACT

Background: 2D Speckle tracking echocardiography (STE) is a non-invasive, angle-independent, semiautomatic and objective technique that quantitatively assesses global and regional longitudinal systolic strain and provides a single bull's eye map for segmental wall strain of the left ventricle. Objectives: assessment of the accuracy of global longitudinal strain (GLS) using STE in the detection of resting myocardial ischaemia and its severity compared with visual assessment of wall motion score index by conventional 2D echo Patients and methods: 100 patients who presented with ACS were included. Wall motion score index (WMSI) was calculated in a 16-segment model and compared with GLS assessed in left ventricle 17 segments and calculated automatically by summation of regional longitudinal peak systolic strain (RLS) using STE. Quantitative coronary angiography was performed on clinical indication and significant stenosis was defined as a 70% reduction of the arterial lumen. Results: 56 patients (56%) of patients were males with a mean age of 58.3 years. GLS showed a significant positive correlation between ejection fraction (EF) and GLS ($p<0.05$), (r value 0.514) and a good significant negative correlation between WMSI and GLS ($p<0.05$), (r value 0.593). And a good significant correlation between GLS and both severity of the lesion and the number of affected vessels was found ($p<0.05$). ROC curves showed the cut-off point of GLS was (-15.9) for predicting lesion $\geq 70\%$ with 88.1% sensitivity, 90.2% specificity respectively, Conclusion: GLS has higher specificity, sensitivity, and diagnostic accuracy for the detection of the severity of lesion and number of the vessel affected than WMSI.