

ملخص البحث الثالث

تاريخ النشر يناير 2022

Comparative Study between Cyanoacrylate Glue versus Sutures for Fixation of Mesh in Inguinal Hernia Open Repair.

Objectives: The Lichtenstein technique is currently the first to repair unilateral primary groin hernias. We aimed to conduct long lateral chain cyanoacrylate as tissue adhesive glue for .open inguinal hernias mesh Fixation

Methodology: 60 patients of inguinal hernia repair were divided randomly into two groups, Group A, sutures did mesh fixation. Group B, mesh fixation was done with cyanoacrylate .glue

Postoperative pain was measured with VAS by direct interview or phone call at 24 hours, 48 hours, seven days, 15 days, one .month, three months, and six months after the operation

Results: There was a statistically significant difference regarding the postoperative pain with a p-value <0.05 between two study groups after 24 hours of operation and after 15 days, 1, 3, and 6 months, with a low mean score among the glue group, which .indicated low pain score among glue group

There was a statistically significant difference with p-value <0.05 between the two study groups regarding operative time with low mean duration among group used glue, which indicated that using glue instead of suture will consume less .time in operation

Conclusion: Mesh fixation with glue causes less postoperative pain, both acute and chronic, than the classical suture fixation, .with similar morbidity and recurrence rates