

Assessment of women Awareness Regard genital prolapse

Mervat Saad Fathy Saad, Dr. Entesar Fatouh Abdel moniem ,

Dr .Rania Abd El- hak Eid Farrag , Dr. Haitham Mohannad Badran

1-Clinical Instructor at Technical Institute of Nursing, Ministry of Health

2-Professor of Maternal and Newborn Health Nursing , Faculty of Nursing - Helwan University

3-Assistant Professor of Maternity and Neonatal Health Nursing , Faculty of Nursing - Fayoum University

4- Assistant Professor of Obstetrics and Gynecology, Faculty of Medicine - Fayoum University

Abstract:

Background:

Genital prolapse is one of the most frequent causes of gynecological morbidities among women in the world. The symptoms of genital prolapse can be extremely debilitating and have an appreciable impact on quality of life aimed to evaluate women awareness regard genital prolapse . A prospective study design was conducted at the out-patient's clinics of Fayoum University hospital and General hospital (obstetrics and gynecology clinic). Purposive sample 73 women was included in the study. Data were collected through three types of tools . It was used to assess genital prolapse stages , Pelvic organ prolapsed-symptoms score (POP-ss) assess symptoms of genital prolapse and . The results of this study revealed that there were a statistically no significant differences between the women with satisfactory level of knowledge and genital prolapsed symptoms, and its stages . While the result Shows that there were statistically non-significant differences between the Total knowledge score and age, residence, income and family member . While the result is almost statistically significant differences between it and level of education, occupation and BMI.

The study recommended increase awareness of women about genital prolapsed symptoms, focusing on issues of outreach and education of the overall population and counseling

program for women with genital prolapse symptoms.

Keywords: prolapse, knowledge , symptoms.