

---

## البحث الخامس

Tamsulosin 0.8 mg daily dose in management of BPH patients with failed tamsulosin 0.4 mg monotherapy and unfit for surgical intervention

جرعة يومية من تامسولوسين 0.8 ملغ في علاج مرضى تضخم البروستاتا الحميد الذين يعانون من فشل العلاج الأحادي تامسولوسين 0.4 ملغ وغير صالح للتدخل الجراحي

(بحث غير مشتق من رسالة)

\* أسم المجلة والناشر:

World journal of Urology - Publisher: Springer.

\* تاريخ النشر:

Accepted: 6/5/2024 Published: 1/6/2024

### English Summary

#### Purpose:

This study aims to evaluate the effectiveness and safety of administering double-dose tamsulosin (0.8 mg) for treating patients with benign prostatic hyperplasia (BPH) who have not responded to the standard single dose of tamsulosin (0.4 mg) and are deemed unsuitable for transurethral resection (TUR) intervention.

#### Patients and Methods:

Between November 2022 and July 2023, we prospectively analyzed 111 patients who were experiencing severe BPH symptoms. These patients received a double dose of tamsulosin for one month. We collected baseline characteristics such as age, body mass index, and underlying medical conditions. Various parameters including the International Prostate Symptom Score (IPSS), prostate-specific antigen (PSA) levels, prostate volume, peak urinary flow rate (Qmax), voided volume, and post-void residual volume were evaluated before and after treatment.

---

### Results:

All 111 patients completed the study. The mean age, PSA level, and prostate volume were  $63.12 \pm 4.83$  years,  $3.42 \pm 0.93$  ng/ml, and  $50.37 \pm 19.23$  ml, respectively. Of these patients, 93 showed improvement in Qmax, post-void residual volume, and IPSS score (p-value = 0.001). The total IPSS score and total Qmax improved from  $24.03 \pm 2.49$  and  $7.72 \pm 1.64$  ml/sec to  $16.41 \pm 3.84$  and  $12.08 \pm 2.37$  ml/sec, respectively.

### Conclusions:

Double-dose 0.8mg tamsulosin as an alpha-blocker therapy appears to be a viable temporary management option for BPH patients who have not responded to the standard single dose 0.4mg tamsulosin and are not suitable candidates for TUR intervention.